



Analyzing the Relationship Between Deforestation and Climate Change Using Satellite Data

Vorobeva Victoria

National Research Tomsk Polytechnic University, Tomsk, Russia

ARTICLE INFO

Keywords:

Deforestation, Climate Change, Satellite Data, Remote Sensing, Greenhouse Gas Emissions

Received: Sep, 23, 2024

Accepted: Nov, 07, 2024

Published: Dec, 25, 2024

ABSTRACT

Deforestation and climate change are two interdependent factors lying at the foundation of global ecosystems and human societies. In this research, the link between climate change and deforestation is searched by employing high-resolution satellite data and advanced analytic methods. Deforestation rates are quantified, changes in land use are assessed, and the emission of greenhouse gases over time is monitored using satellite imagery and remote sensing data. Also, the study examines the thermal variations and precipitation patterns on a regional and global scale as well as carbon recruitment capabilities in the affected areas. Moreover, this research paper makes a thorough analysis of how deforestation activities cause global climate changes by integrating different metrics derived from satellite data and climate models. The results underline the importance of forest conservation in the fight against climate change and put forward practical proposals for the development of policies and land use sustainable practices. The ultimate objective of this study is to create a better understanding of the complex interactions between climate change and deforestation, thus creating a basis for addressing in a data-driven way such challenges as the ones that affect our planet on a global level.

1. INTRODUCTION

Environmental issues like deforestation are now the primary cause of climate change. The enormous forest devastation is indeed the cause of many life-threatening drawbacks, bringing about far-reaching effects which alter ecosystems, affect biodiversity, and cause climate systems to be unpredictable (Choudhary et al., 2023). Timberlands act as a key part of the Carbon cycle by acting as massive sinks of carbon dioxide (CO₂) and also serve the function of detoxifying the Earth's atmosphere. Deforestation, on the other hand, is disrupting this balance, venting the stored carbon back into the atmosphere and leading to global warming (Kaçmaz Akkurt, 2023). Although people have voiced concern and put in place policies to address this issue in many parts of the

world for many decades, it is now even far worse in some areas, as agricultural expansion, urbanization, logging, and other human activities are the primary reasons for the fast deforestation (Gomathy & Kalaiselvi, 2023). Therefore, the intersection of deforestation and climate change is a very complex and multifaceted issue that necessitates rigorous and nuanced data-driven exploration if we are to mitigate it effectively and manage it sustainably (Guo et al., 2021).

Thanks to satellite technology and remote sensing, we can now monitor and analyze environmental changes on a global scale like never before. Until now, deforestation studies have relied on ground-based surveys and regional data that are often limited in scope and therefore widely unreliable. By contrast, satellite data offers a far more

comprehensive perspective; high-resolution imagery and the temporal coverage provided by the satellites enable researchers to monitor deforestation in real time and over longer periods of time (Hasegawa et al., 2022). Remote sensing technology can not only notice any slightest shifts in land cover but also quantify the loss of forests and measure exactly the most important variables like the condition of the vegetation, density of the canopy, and emissions of carbon. These abilities make it impossible to ignore satellite data when discussing the relation between deforestation and climate change. Moreover, combining satellite data with climate models, increases the capability of predicting trends in the future and assessing the impacts of forestry on the world, as well as regional climate systems (Abramoff et al., 2023).

The relationship between deforestation and climate change is complex and involves various direct and indirect mechanisms. Carbon dioxide and other greenhouse gases are directly emitted when trees and other plants are removed or burnt, contributing to global warming. Trees are the best natural absorbers of carbon dioxide as they utilize it for growth through photosynthesis. Carbon dioxide that was previously stored in the trees and soil is released into the atmosphere when forests are cleared or burnt, leading to the rise of greenhouse gases. It is estimated that deforestation accounts for approximately 10% of greenhouse gas emissions worldwide each year. In an indirect manner, deforestation influences local and regional wildlife diversity by changing the surface albedo, reducing evapotranspiration, and affecting the precipitation patterns (Crane-Droesch, 2018). Consequently, these effects can cause an increase in temperature, altered weather systems, and long-term climate instability. Furthermore, deforestation reduces the diversity of plants and animals, thus making ecosystems less capable of overcoming environmental stressors like climate change and makes them more prone to adaptation challenges (Sidhu et al., 2023).

It is important to note that deforestation has implications that go beyond climate systems and involve social, economic, and ecological factors. For instance, vital tropical rainforests are the Amazon, Congo Basin, and those in Southeast Asia in supporting indigenous communities, providing ecosystem services, and conserving global biodiversity. The loss of forests in these areas

compromises the livelihoods of millions of people, disrupts the water cycles, and threatens forest-dependent species (Gyamerah et al., 2023). Furthermore, the reduction of forest cover can cause several problems in agriculture, water availability, and human health, creating feedback loops that enhance climate vulnerabilities. The understanding of such connected ramifications is essential for developing comprehensive solutions that can handle both deforestation as well as its far-reaching consequences (Ravindiran et al., 2023). Understanding the relationship between deforestation and climate change is essential for the successful implementation of global environmental initiatives. By way of example, the Paris Agreement and the United Nations Sustainable Development Goals (SDGs) continue to stress the imperative of combating deforestation within climate action programs. Nevertheless, the successful execution of such programs should be based on the gathering of solid data and scientific acumen. Satellite data stands out as a significant opportunity to overcome the knowledge gap, as it presents refined and realistic information on deforestation trends, hot spots, and corresponding climatic ramifications. Such a data-driven view is capable of informing policy decisions, specifying intervention priorities, and supporting worldwide action against climate change (Li et al., 2023).

In spite of the rising availability of satellite data, there are still obstacles to the investigation and interpretation of the complex relationships between deforestation and climate change. The diversity of drivers of deforestation, regional discrepancies, and the fact that they have non-linear feedback mechanisms, are significant barriers to analysis. Moreover, the effects of deforestation on the climate may not be the same in all areas but depend on the geographical ones, forest types, and of course on the scale of land-use change (Ramachandra et al., 2023). For instance, while tropical forests, which contain large amounts of carbon and influence global weather patterns, show different climate medications as opposed to boreal or temperate forests, the socio-economic conditions, governance systems, and law enforcement practices shape the outcomes of deforestation decisions, thus making the analysis even more complicated. To overcome these difficulties, it is necessary to engage in an interdisciplinary approach that would combine

satellite data, ground-based observations, climate models as well as social-economic knowledge (Antonelli et al., 2023).

In this research, we intend to employ satellite data to exhaustively examine the interconnection between deforestation and climate change. The integration of remote sensing technologies along with sophisticated analytical models is intended to allow us to compute the level and pace of deforestation, attribute it to greenhouse gas emissions, and perceive its impact on the local and global climate systems. We emphasize identifying trends, correlations, and causal mechanisms that can guide the policy and conservation decisions. Hence, the research sought to address the following inquiries (McCallum et al., 2023): How does deforestation augment the greenhouse gas emissions? What are the regional and global climate outcomes due to the disappearance of forest cover? And to what extent can satellite data enhance the monitoring and control strategies?

Objectives

- The application of satellite data and remote sensing methods helps quantify the emissions of greenhouse gases.
- A careful examination of the roles of deforestation in both the regional and global climate, covering such aspects as carbon cycle, precipitation, and temperature, would be performed.

Once these objectives are met, this research aims to deepen the knowledge of the impact of deforestation on climate change and assist those who support forest conservation based on scientific evidence. The results of this study will be relevant to government leaders, environmental conservators, and scientists as they tackle both the issues of deforestation and climate change. In addition, our method shows that satellite data can be an essential element in exactly, successfully, and repeatedly monitoring the environment. Thus, it may lead to different ways of studying climate and advancement in sustainability .

2. LITERATURE REVIEW

The quest to comprehend the multi-layered interdependencies between forest desolation and climate evolution calls for a thorough examination of precedently published reports. Various investigations have illicitly discussed the reason, consequence, and transition strategies topical to

forest degradation but only a few of them have highlighted the inextricable relationship of global warming, local and universal climate shifts, and the scientific instruments of the technology segment for monitoring and analyses. The linkage of satellite and remote sensors to researchers left them to produce a plethora of details, documenting deforestation processes and their climate repercussions. In this part, the crux of the significant studies that led to the consummate knowledge of how deforestation affects climate systems will be summarized along with the procedures, the results, and the shortcomings of old studies (Altarez et al., 2023).

(Sboui et al., 2023) thoroughly investigated how machine learning can be used to improve the forecast of deforestation associated with oil palm plantations. The research proposed a novel conceptual framework that merges geospatial technologies with image processing techniques to confront the menace of palm oil-related deforestation, which has often been neglected in the current literature. The framework was composed of three parts: data preparation, model training, and validation, in Aceh region of Indonesia (Duku & Hein, 2023). Results obtained from their sensitivity analysis shows that the accuracy of the deforestation rates prediction with the proposed approach can be achieved. This study points out that machine learning plays an essential role in the support of decision-makers through offering of actionable insights on the possibility of where and when the new plantations could be established sustainably. This research stresses the usability of the concept of predictive analytics in the management of deforestation for the purpose of environmental damage control.

Daiyoub et al. , in their study of the forests of Syria, looked into the effects of armed conflict on the forest trees, along with other forest lands between 2010 and 2019 (Daiyoub et al., 2023). By employing Landsat and PALSAR satellite images, as well as the Tasseled Cap transformation and the LandTrendr algorithm, the study showed a significant reduction of 19.3% of the forest cover. Among the most important factors were illegal logging close to the refugee camps, the proximity of roads and settlements, as well as the forest fires resulting from explosive events, which were the bombings. The authors emphasized on finding sustainable alternatives for heating and cooking to

avoid the dependency on deforestation during the post conflict reconstruction process. The research presents persuasive evidence showing that the socio-political factors affect the deforestation process, and the needs for better governance and enforcement of policies are vital for forest conservation (Yuh et al., 2023).

Silva et al. analyzed the relationship between management practices and deforestation in Brazil's 283 protected areas (PAs) (da Silva et al., 2023). The data concluded that the machine was quite good at identifying ones with a risk of deforestation and wildfires with accuracy rates of 87.5% and 72.8%, respectively. It was identified that biome, area, tourism management, and governance were the critical factors that should be taken into consideration in predicting vulnerability. The sustainable management of tourism was said to be the best way of minimizing deforestation and increasing the effectiveness of the governance of the protected area. Herewith the interlinking of tourism and conservation efforts is highlighted and therefore it is recommended for policies to be made that benefit from the tourism sector and thus preserve the resources of the protected areas.

(Butt et al., 2023) studied the effects of tropical deforestation on local and regional climate warming, focusing on the Amazon rainforest. The study, with the use of the remote-sensing observations and machine learning showed that Deforestation caused warming even at distances up to 100 km from forest loss sites. The researchers concluded that the loss of 10% of the forest would

result in a four-fold increase in the temperature and that the weather resilience of the regions would be in danger (Frimpong et al., 2023). The researchers made a forecast, that if the deforestation process continues, the dry season warming in the south of the country may reach 0.96 K by 2050, while in the case of the elimination of the deforestation process risk it may reduce the warming up to 0.4 K. The study is of great importance because it shows the need for avoiding deforestation, so that the climate regions can be made less vulnerable and the sustainability of the tropical regions can be enhanced.

Picanço Rodrigues and Caetano explored the influence of political activity on deforestation and fire outbreaks in the Brazilian Amazon through the application of sentiment analysis derived from Twitter data (Picanço Rodrigues & Leonel Caetano, 2023). The analysis of satellite fire data as well as social media engagement determined that positive comments in respect of agriculture and deforestation-related policies usually came six days before the fire breakout. The decision tree regressor was identified as the most potent predictive model. The outcomes of this research are a direct reflection of the influence political stories have on educating and changing the public's attitudes toward the destruction of the environment. This study implies the use of machine learning and sentiment analysis over human-induced crises as well as a need for stricter policies to be adopted in political contexts where deforestation is concerned.

Table 1 : Comparson of the Related Studies

Study	Focus Area	Methodology	Key Findings	Applications
Sboui et al. (2023)	Deforestation due to oil palm plantations	Machine learning framework using geospatial data	Developed a predictive framework for deforestation due to oil palm, achieving accurate predictions in the Aceh province of Indonesia. Sensitivity analysis validated the framework's effectiveness.	Helps decision-makers manage palm oil-related deforestation sustainably.
Daiyoub et al. (2023)	Impact of armed conflict on forest cover	Remote sensing with Landsat and PALSAR; LandTrendr	Identified a 19.3% forest cover loss in Syria (2010-2019) caused by illegal logging, proximity to refugee camps, roads, and bombings. Emphasized post-conflict sustainable alternatives for fuel to mitigate deforestation.	Informing post-conflict forest restoration and sustainable resource management strategies.
Silva et al. (2023)	Protected area management	Machine learning classification and	Identified tourism management and governance as key	Developing policies for sustainable tourism to

	and deforestation	multivariate models	predictors of deforestation vulnerability in Brazil's protected areas. Highlighted tourism's potential role in reducing deforestation and enhancing governance effectiveness.	reduce deforestation and improve protected area management.
Butt et al. (2023)	Regional climate impacts of deforestation	Remote sensing and machine learning	Showed that deforestation in the Amazon caused warming up to 100 km away. Nonlocal warming increased by a factor of four due to deforestation. Projected future warming scenarios under different deforestation trajectories.	Strengthening regional climate adaptation and resilience strategies through reduced deforestation.
Picanço Rodrigues & Caetano (2023)	Political narratives and deforestation	Machine learning and sentiment analysis of social media	Found political narratives influence fire outbreaks and deforestation in the Brazilian Amazon. Positive comments on deforestation policies were predictive of fire events, with a 6-day lag identified via decision tree regression.	Utilizing social media analysis to detect and mitigate politically-driven deforestation trends.

3. PROPOSED METHODOLOGY

This investigation uses an all-encompassing methodology that includes satellite data, sophisticated remote sensing techniques, and machine learning models for examining the correlation between deforestation and climate change. The primary objective is to utilize the latest technologies to gather, process, and analyze high-resolution data to get practical insights into the forest alteration dynamics and their effect on the global climate. The stratagem is classified into various linked steps like the acquisition of satellite data, data preprocessing, metadata extraction, storing, analysis, and publication. By designing these modes, we ensure the precise, replicated, and systematic assessment of deforestation and its climatic consequences.

The initial phase of the methodology comprises the acquisition of satellite data. High-resolution imagery was obtained from diverse sources like satellites with special sensors capable of taking Normalized Difference Vegetation Index (NDVI), thermal imaging, and atmospheric CO₂ readings. NDVI serves as a vital marker of the health of vegetation and thus called to distinguish between areas under dense forest cover, degraded zones, and non-vegetated regions. Furthermore, the variant of thermal sensors was used in the detection of changes in surface-land temperatures,

which are very often linked to deforestation. Other sensors, on the other hand, are used to monitor the emissions of greenhouse gases, especially CO₂, released through deforestation. Consequently, these data sources together become the strongest basis for the assessment of forest destruction and its effect on climate.

The preprocessing of the raw satellite data happens after data acquisition. The data is then conveyed to ground stations where telemetry analysis, and preliminary filtering are carried out to get rid of noise and ensure reliability via the use of a removal filter. Data preprocessing entails atmospheric correction to prevent the distortion caused by the atmospheric particles, cloud removal to obtain clarity in the data, and cleaning of the data to get rid of the inconsistencies. These bettered procedures will verify the credibility of the performance of the analysis. In addition, metadata with timestamps and geospatial coordinates will be extracted for the temporal analysis as well as the spatial analysis. The formatting of raw data was the preprocessing stage of it is the gateway to further data processing ones and consequently, the analysis.




Figure 1 : Working of the Proposed Model
 The processing of the satellite data is followed by storing it in structured databases, by utilizing cloud-based platforms like AWS and Google Cloud, the organization, and indexing of the satellite data will be possible. These will, in turn, ensure the

speedy and efficient retrieval of the data which is subsequently to be analyzed. The databases are not only capable of processing the time and space data but also they can do research that tracks deforestation patterns over time and studies them with respect to the environment. With the use of cloud storage solutions, we have scalability as a quick solution for handling a lot of data since we can just integrate new data as it becomes available. The platforms also give us computational strength which is useful for running complicated analytical models on large datasets.

The methodology is culminating in the data analysis. The techniques for feature extraction are utilized to come up with useful variables such as vegetation indices, the rates of deforestation, and climate indicators. Then, the machine learning models are used to analyze these features over time and figure out the trends, correlations, and patterns. Thus, predictive models will provide the estimated future deforestation rates and the possible effects on climate systems. The relation of temperature, precipitation, and carbon emissions to the forest, in effect, will be regarded as deforestation through the correlation method. Long-term changes will be detected by using the trend prediction model. The implication of the availability of the machine learning would be that the complex interactions that are not easily discerned in the method can be detected.

Visualization tools are used to present the results in a way that is intuitive and easily accessible. Geospatial heatmaps indicate hotspots of deforestation, while temporal charts exemplify changes over time. Advanced 3D renderings provide a deeper immersion, allowing stakeholders to see the magnitude of forest loss and its climatic effects without complex textual data. The role of these visualizations in the communication of the research results to policymakers, environmental organizations, and the general public is key. The transformation of complex data into visually appealing graphics is the method by which the research tries to overcome the gap between scientific evidence and achievable actions.

The methodology's last step consists of putting the results together in comprehensive reports and direct actions. The data analysis results are used as the basis for policy suggestions, outlining environmental risks, and shooting eligibility

criteria. These reports are written to meet the distinct needs of the various stakeholders, such as the NGOs. Access to datasets and statistical tools of high quality can enable the research community to undertake further studies. The policymakers get solid proof of the recommendations that will be the basis of the legislative structure and climate action plans. The NGOs have the information they need to raise funds for conservation and mobilization programs. The research wants to serve the various capabilities of the stakeholders so that the greater device can be reached in the fight against deforestation and climate change.

The functionality of the model suggested is as illustrated in Figure 1. The model signifies a systematic framework for the gathering, processing, and analysis of satellite data along with multiple technological aspects whose integration is ensured to generate results that are both accurate and comprehensive. A satellite is used to collect data, such as detailed images, NDVI metrics, and atmospheric indicators. After this, the information is sent to ground stations by the telemetry analysis and initial processing. Afterward, the refined data is indexed and kept in the cloud-based repositories guaranteeing accessibility and flexibility of resourceful retrieval. Data interpretation is then done using sophisticated analytical techniques including the implementation of machine learning models. As a visualization illustration of these results, geospatial heat maps, temporal graphs, and 3D renderings are utilized. The results obtained from this model are gradually combined into comprehensive reports that offer actionable recommendations for decision-makers, researchers, and NGOs.

The model outlined above is an exemplary representation of the coupling of satellite technology with advanced analytics to create durable solutions for environmental issues. This model aims at a comprehensive approach to monitor and analyze both deforestation and climate change; thus, it is a contribution to the formation of sustainable strategies. Also, it highlights the need to utilize technology appropriately for better quality, quantity, and effect in environmental research.

4. SIMULATION RESULTS

This research's conclusions give imperative revelations into the link between

deforestation, carbon discharges, and climate change. A Kaggle centered dataset, blended with the relevant UCI land cover data information, was chiefly used for the analysis. Both tables and various graphs were used in reports respectively to illustrate trends and correlations with the periods and specific variables, such as deforestation rates, CO₂ emissions, and global temperature anomalies. The descriptive statistics of the datasets show major differences in deforestation rates, with an average annual decrease of 0.998 million hectares between 2000 and 2022. The average amount of carbon dioxide emissions was 3.24 billion metric tons per year, which remarkably indicates the role of deforestation in the alteration of the concentration of greenhouse gases in the atmosphere. The global temperature rise, which is a good indicator of climate change, showed an average growth of 1.02°C in the compared reference period while the pre-industrial one was used to estimate it, emphasizing the global warming event as a climatic result of deforestation and logging activities.




Figure 2 : Deforestation Rate Over Time (2000-2022)




Figure 3 : CO2 Emissions Over Time (2000-2022)




Figure 4 : Global Temperature Anomaly Over Time (2000-2022)

In a more visual way, these trends can be further revealed through graphical representations. Figure 2 displays the deforestation rate as a function of time, showing that the total number of trees had a tendency to oscillate, sometimes there was an increase in forest loss during the time when agricultural activity and industry expanded quite quickly. This phenomenon is like the figures of 2010 and 2020; the former shows a striking peak in logging and the latter one, widespread land clearing, which probably caused the economic pressure and so this was the first one of the important signals of the unsustainable management of forests.

In the Figure 3, CO₂ emissions throughout the same

period and the deforestation rates show a very strong positive correlation. The emissions rise in peak years of deforestation in the same way as they confirm that the removal of trees from the forests is the process that releases greenhouse gases into the atmosphere. The trends in emissions and deforestation measured in that way would mitigate greenhouse gas emissions.

Figure 4 illustrates the global temperature anomaly, which has steadily increased over the years. Although temperature variations are influenced by several factors, the observed trend matches the rising carbon emissions due to deforestation. The data indicate that deforestation not only leads to immediate carbon emissions but also disrupts climate stability in the region and so it leads to environmental degradation globally.

These findings are consistent with the existing literature that explains the dual role of forests in carbon sequestration and climate regulation. This research had employed a direct combination of satellite data with Machine Learning models enabling the detection of the mapping of complex relationships between deforestation and climate change beyond the previous research experience. The graphical outputs and summary statistics provide a robust basis for understanding the temporal and spatial dynamics of these interactions.

Table 2 : Summary Statistics Of Deforestation And Climate Change Data

	Count	Mean	Std Dev	Min	25%	Median	75%	Max
Year	23	2011.00	6.78	2000.00	2005.50	2011.00	2016.50	2022.00
Deforestation Rate (Million ha)	23	0.998	0.291	0.596	0.696	1.028	1.208	1.451
CO ₂ Emissions (Billion tons)	23	3.238	0.661	2.291	2.730	2.993	3.817	4.302
Temperature Anomaly (°C)	23	1.023	0.120	0.818	0.907	1.049	1.119	1.180

As depicted in the proposed model (Figure 1 in the Methodology section), the integrated analysis framework used in this study emphasizes the rationale for using satellite data in environmental monitoring. The method not only records the number of trees that have been lost but also it investigates the climatic ramifications of the deforestation, thus providing a certain approach which is replicable for the future studies of this

kind. Moreover, the detailed results give some direction to the studies for the possible actions that would be taken up by the different groups of people, such as governments, scientists, and environmental organizations, in the fight against the two crises of deforestation and global warming.

5. CONCLUSION

This study comprehensively examines the significant relationship between deforestation and

climate change by utilizing an analysis of the satellite data which is the source of information about forest loss, CO₂ emissions, and global temperature anomalies. During the period of 2000-2022, an average annual deforestation rate of 0.998 million hectares was stated as a fact, which caused a sufficient amount of CO₂ emissions to occur in the atmosphere through an average of 3.24 billion metric tons each year. Besides, the analysis of the results indicated, a steady increment in the rate of global temperature anomalies was observed, which was equal to the rate of warming of 1.02 °C above pre-industrial levels, illustrating that the impact of deforestation on global warming was seminal. Graphical displays confirmed that trees of high deforestation were the principal and basic correlations of the peaks in CO₂ emissions, so there is an urgent need for effective developmental forest conservation actions to alleviate climate changes.

Despite the useful information being provided by this research, some limitations cannot be ignored. Trusted by global satellites, the analysis may have the natural biases of cloud cover or effects in the atmosphere due to atmospheric disturbances which are despite the preprocessing work of the data. Also, the study's excluding of the regional socio-economic drivers of deforestation is responsible for missing the additional context that complements the findings. For a better understanding of the implications of deforestation and to explore advanced machine learning models for predictive accuracy, future research must focus on bringing together localized data and economic variables within the study. The core message remains that this study reiterates the fact that forests are a key aspect of the regulation of the climate, while it also highlights the necessity of using data-driven interventions to tackle deforestation and climate change.

REFERENCES

Abramoff, R. Z., Ciaias, P., Zhu, P., Hasegawa, T., Wakatsuki, H., & Makowski, D. (2023). Adaptation strategies strongly reduce the future impacts of climate change on simulated crop yields. *Earth's Future*, 11(4), e2022EF003190. <https://doi.org/10.1029/2022EF003190>

Altarez, R. D. D., Apan, A., & Maraseni, T. (2023). Deep learning U-Net classification of Sentinel-1 and 2 fusions effectively demarcates tropical montane forest's deforestation. *Remote Sensing Applications: Society and Environment*,

29, 100887. <https://doi.org/10.1016/j.rsase.2022.100887>

Antonelli, A., Dhanjal-Adams, K. L., & Silvestro, D. (2023). Integrating machine learning, remote sensing and citizen science to create an early warning system for biodiversity. *Plants, People, Planet*, 5(3), 305–316. <https://doi.org/10.1002/ppp3.10337>

Antonelli, A., Dhanjal-Adams, K. L., & Silvestro, D. (2023). Integrating machine learning, remote sensing and citizen science to create an early warning system for biodiversity. *Plants, People, Planet*, 5(3), 305–316. <https://doi.org/10.1002/ppp3.10337>

Butt, E. W., Baker, J. C. A., Silva Bezerra, F. G., von Randow, C., Aguiar, A. P. D., & Spracklen, D. V. (2023). Amazon deforestation causes strong regional warming. *Proceedings of the National Academy of Sciences*, 120(45), e2309123120. <https://doi.org/10.1073/pnas.2309123120>.

Choudhary, P., Shukla, P., & Muthamilarasan, M. (2023). Genetic enhancement of climate-resilient traits in small millets: A review. *Heliyon*, e14502. <https://doi.org/10.1016/j.heliyon.2023.e14502>

Crane-Droesch, A. (2018). Machine learning methods for crop yield prediction and climate change impact assessment in agriculture. *Environmental Research Letters*, 13(11), 114003. <https://doi.org/10.1088/1748-9326/aae159>

da Silva, F. R., et al. (2023). Machine learning application to assess deforestation and wildfire levels in protected areas with tourism management. *Journal for Nature Conservation*, 74, 126435. <https://doi.org/10.1016/j.jnc.2023.126435>

Daiyoub, A., Gelabert, P., Saura-Mas, S., & Vega-Garcia, C. (2023). War and deforestation: Using remote sensing and machine learning to identify the war-induced deforestation in Syria 2010–2019. *Land*, 12(8), 1509. <https://doi.org/10.3390/land12081509>

Duku, C., & Hein, L. (2023). Assessing the impacts of past and ongoing deforestation on rainfall patterns in South America. *Global Change Biology*, 29(18), 5123–5138. <https://doi.org/10.1111/gcb.16856>

Frimpong, B. F., Koranteng, A., Atta-Darkwa, T., Junior, O. F., & Zawiła-Niedźwiecki, T. (2023). Land cover changes utilising Landsat satellite imageries for the Kumasi Metropolis and its adjoining municipalities in Ghana (1986–2022). *Sensors*, 23(5), 2644. <https://doi.org/10.3390/s23052644>

Gomathy, M., & Kalaiselvi, K. (2023). Climate change and its impact on agriculture. In *Advanced technologies for smart agriculture*. *International Journal of Zoology and Applied Biosciences*. <https://doi.org/10.55126/ijzab.2024.v09.i04.004>

Guo, Y., et al. (2021). Integrated phenology and climate in rice yields prediction using machine learning methods. *Ecological Indicators*, 120, 106935. <https://doi.org/10.1016/j.ecolind.2020.106935>

Gyamerah, S. A., Asare, C., Mintah, D., Appiah, B., & Kayode, F. A. (2023). Exploring the optimal climate conditions for maximum maize production in Ghana: Implications for food security. *Smart Agricultural Technology*, 6, 100370. <https://doi.org/10.1016/j.jatech.2023.100370>

Hasegawa, T., et al. (2022). A global dataset for the projected impacts of climate change on four major crops. *Scientific*

- Data, 9(1), 150. <https://doi.org/10.1038/s41597-022-01150-7>
- Hu, T., et al. (2023). Crop yield prediction via explainable AI and interpretable machine learning: Dangers of black box models for evaluating climate change impacts on crop yield. *Agricultural and Forest Meteorology*, 336, 109458. <https://doi.org/10.1016/j.agrformet.2023.109458>
- Kaçmaz Akkurt, G. (2023). Vulnerability to climate change. In *Sustainability, conservation and ecology in spatial planning and design: New approaches, solutions, applications*. IGI Global. <https://doi.org/10.4018/978-1-5225-0803-8.ch066>
- Li, L., et al. (2023). Integrating machine learning and environmental variables to constrain uncertainty in crop yield change projections under climate change. *European Journal of Agronomy*, 149, 126917. <https://doi.org/10.1016/j.eja.2023.126917>
- McCallum, I., et al. (2023). Crowd-driven deep learning tracks Amazon deforestation. *Remote Sensing*, 15(21), 5204. <https://doi.org/10.3390/rs15215204>
- Picanço Rodrigues, V., & Leonel Caetano, M. A. (2023). The impacts of political activity on fires and deforestation in the Brazilian Amazon rainforest: An analysis of social media and satellite data. *Heliyon*, 9(12), e22670. <https://doi.org/10.1016/j.heliyon.2023.e22670>
- Pisl, J., Hughes, L. H., Ruswurm, M., & Tuia, D. (2023). Classification of tropical deforestation drivers with machine learning and satellite image time series. In *Proceedings of the International Geoscience and Remote Sensing Symposium (IGARSS)*. IEEE. <https://doi.org/10.1109/IGARSS52108.2023.10281472>
- Ramachandra, T. V., Mondal, T., & Setturu, B. (2023). Relative performance evaluation of machine learning algorithms for land use classification using multispectral moderate resolution data. *SN Applied Sciences*, 5(10), 1546. <https://doi.org/10.1007/s42452-023-05496-4>
- Ravindiran, G., et al. (2023). Impact of air pollutants on climate change and prediction of air quality index using machine learning models. *Environmental Research*, 239, 117354. <https://doi.org/10.1016/j.envres.2023.117354>
- Sboui, T., Saidi, S., & Lakti, A. (2023). A machine-learning-based approach to predict deforestation related to oil palm: Conceptual framework and experimental evaluation. *Applied Sciences*, 13(3), 1772. <https://doi.org/10.3390/app13031772>
- Sidhu, B. S., Mehrabi, Z., Ramankutty, N., & Kandlikar, M. (2023). How can machine learning help in understanding the impact of climate change on crop yields? *Environmental Research Letters*, 18(2), 024027. <https://doi.org/10.1088/1748-9326/acb164>
- United Nations. (n.d.). Take urgent action to combat climate change and its impacts. United Nations. <https://sdgs.un.org/goals/goal13>
- Yuh, Y. G., Tracz, W., Matthews, H. D., & Turner, S. E. (2023). Application of machine learning approaches for land cover monitoring in northern Cameroon. *Ecological Informatics*, 74, 101955. <https://doi.org/10.1016/j.ecoinf.2022.101955>